

(12) UK Patent Application (19) GB (11) 2 106 902 A

(21) Application No 8227449

(22) Date of filing
27 Sep 1982

(30) Priority data

(31) 3138550

(32) 28 Sep 1981

(33) Fed Rep of Germany
(DE)

(43) Application published
20 Apr 1983

(51) INT CL³ C07D 213/26

A61K 31/44

C07D 401/12

(C07D 213/26 207/00

213/00 295/00)

(52) Domestic classification

C2C 1530 215 220 22Y

250 251 25Y 29X 29Y

30Y 313 314 21Y 321

322 323 32Y 332 337

338 364 365 36Y 373

37Y 456 45Y 493 500

50Y 614 620 621 623

624 630 634 644 650

660 661 662 670 672

680 682 69Y 758 779

802 80Y AA LF LH LW

NG RM

U1S 1327 C2C

(56) Documents cited

None

(58) Field of search

C2C

(71) Applicant

Boehringer Ingelheim

International GmbH

(FR Germany)

D-6507

Ingelheim/Rhein

Federal Republic of

Germany

(72) Inventors

Claus Schneider

Gerhard Walther

Wolf Dietrich Bechtel

Karl-Heinz Weber

Karin Boke-Kuhn

(74) Agents

Frank B Dehn and Co


Imperial House

15-19 Kingsway

London WC2B 6UZ

(54) 2,2-Disubstituted-ethylamines

(57) Substituted 2-phenyl-2-(pyridyloxy)-ethylamines and isosteric compounds of general formula



wherein

R₁ and R₂, which may be the same or different, each represents a hydrogen or halogen atom or a methyl, methoxy, amino or nitro

group;

R₃ represents a hydrogen or halogen atom or a methyl group;

R₄ and R₅, which may be the same or different, either each represents a hydrogen atom or an alkyl group with 1 or 2 carbon atoms or, together with the nitrogen atom to which they are attached they may form a pyrrolidino or morpholino ring, and

X represents O, HN or S;

and acid addition salts thereof.

The new compounds are highly effective antidepressants with a structure hitherto unknown in this field of application.

GB 2 106 902 A


SPECIFICATION

2,2-Disubstituted-ethylamines

5 The invention relates to new substituted 2-phenyl-2-(pyridyloxy)-ethylamines and isosteric compounds, to processes for their preparation, to pharmaceutical compositions containing them and to their use as antidepressants.

According to one feature of the present invention there are provided substituted 2-phenyl-2-(pyridyloxy)-ethylamines and isosteric compounds of general formula

10



wherein

25 R_1 and R_2 , which may be the same or different, each represents a hydrogen or halogen atom or a methyl, methoxy, amino or nitro group;

R_3 represents a hydrogen or halogen atom or a methyl group;

R_4 and R_5 , which may be the same or different, either each represents a hydrogen atom or an alkyl group with 1 or 2 carbon atoms or, together with the nitrogen atom to which they are


30 attached they may form a pyrrolidino or morpholino ring, and

X represents O, HN or S;

and acid addition salts thereof.

These compounds according to the invention possess interesting pharmacological properties and in particular a useful antidepressant activity. It will be appreciated that, for pharmaceutical use, the salt referred to above will be physiologically compatible but other salts may find use, for example, in the preparation of compounds of general formula I and their physiologically compatible salt.

35 The new compounds may be obtained by reacting a phenylethylamine of formula



(wherein R_1 , R_2 , R_4 and R_5 are as hereinbefore defined and Y represents a halogen atom or a hydroxy group), or an acid addition salt of this compound, with a pyridine derivative of formula



55 (wherein R_3 is as hereinbefore defined and Z represents a hydroxy, amino or mercapto group) followed, if desired, by conversion of the product thus obtained into an acid addition salt thereof.

In the case of starting materials of general formula II wherein Y represents a hydroxy group, condensation is preferably effected with a compound of formula III in an acidic medium at elevated temperatures. For this reaction, it is possible to use strong inorganic or organic acids, e.g. concentrated mineral acids such as for example hydrochloric acid, hydrobromic acid, sulphuric acid, or polyphosphoric acid and also acetic anhydride, phosphorus pentoxide or trifluoroacetic acid.

60 Phenylethylamines of general formula II wherein Y represents a halogen atom are preferably reacted with the pyridine derivative of formula III in the presence of a base and preferably under

65

phase transfer conditions (i.e. with the addition of one or more long-chained tertiary amines such as e.g. a benzyl-tributyl-ammonium halide or a tetrabutyl-ammonium halide or using benzyl-triphenyl-phosphonium chloride).

Suitable bases include, for example, inorganic bases such as e.g. alkali and alkaline earth metal hydroxides and carbonates and organic bases such as e.g. pyridine, piperidine, tertiary amines and alkali metal alkoxides.

Compounds according to the invention wherein R_4 and/or R_5 represent hydrogen may be alkylated in a manner known *per se*, e.g. by reacting them with a dialkyl sulphate or an alkyl halide, to give the corresponding compound wherein R_4 and/or R_5 represents an alkyl group with 1 or 2 carbon atoms.

Using the process described above, the following end products may, for example, be obtained optionally in the form of their acid addition salts:

- 2-phenyl-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine, 15
- 2-phenyl-2-(pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine, 15
- 2-(p-bromophenyl)-2-(pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine;
- 2-(p-bromophenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine;
- 2-(p-bromophenyl)-2-(2-chloropyridyl-(3)-oxy)-N,N-dimethyl-ethylamine,
- 2-(p-bromophenyl)-2-(5-chloro-pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine,
- 20 2-(p-bromophenyl)-2-(6-chloro-pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine, 20
- 2-(p-bromophenyl)-2-(pyridyl-(4)-oxy)-N,N-dimethyl-ethylamine,
- 2-(p-chlorophenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine,
- 2-(m,p-dichlorophenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine,
- 2-phenyl-2-(pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
- 25 2-phenyl-2-(pyridyl-(3)-amino)-N,N-dimethyl-ethylamine, 25
- 2-(p-bromophenyl)-2-(pyridyl-(2)-amino)-N-methyl-ethylamine,
- 2-(p-bromophenyl)-2-(pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
- 2-(p-bromophenyl)-2-(pyridyl-(3)-amino)-N,N-dimethyl-ethylamine,
- 2-phenyl-2-(4-methyl-pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
- 30 2-(p-bromophenyl)-2-(5-methyl-pyridyl-(2)-amino)-N,N-dimethyl-ethylamine, 30
- 2-(p-bromophenyl)-2-(6-methyl-pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
- 2-(4-p-bromophenyl)-2-(5-chloro-pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
- 2-(4-p-bromophenyl)-2-(pyridyl-(2)-thio)-N,N-dimethyl-ethylamine,
- 2-(p-bromophenyl)-2-(pyridyl-(3)-oxy)-morpholino-ethylamine,
- 35 2-(p-bromophenyl)-2-(pyridyl-(3)-oxy)-pyrrolidino-ethylamine, 35
- 2-(p-bromophenyl)-2-(pyridyl-(2)-amino)-morpholino-ethylamine,
- 2-(p-bromophenyl)-2-(pyridyl-(2)-amino)-pyrrolidino-ethylamine,
- 2-(p-methoxyphenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine,
- 2-(p-hydroxyphenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine,
- 40 2-(p-tolyl)-2-(pyridyl-(3)-oxy)-N,N-diethyl-ethylamine, 40
- 2-(p-nitrophenyl)-2-(pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine,
- 2-(4-aminophenyl)-2-(pyridyl-(2)-oxy)-N,N-dimethyl-ethylamine,
- 2-(m,p-dichlorophenyl)-2-(pyridyl-(3)-amino)-N,N-dimethyl-ethylamine,
- 2-(m,p-dimethoxyphenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine.

45 The starting materials are widely used chemicals which may be obtained commercially or can be prepared using methods which are generally known. 45

The end products of general formula I may, if desired, be converted into the acid addition salts thereof by conventional methods.

Suitable acids include both inorganic acids such as e.g. hydrohalic acids, sulphuric, phosphoric and amino-sulphonic acid, and also organic acids such as e.g. formic, acetic, propionic, lactic, glycolic, gluconic, maleic, succinic, tartaric, benzoic, salicylic, citric, ascorbic, p-toluenesulphonic and oxyethane sulphonc acid. 50

As mentioned above the compounds of general formula I and the acid addition salts thereof possess interesting pharmacological properties. Those which we have tested have proved highly effective in biochemical and pharmacological test systems specific to antidepressants. Thus, they are capable of inhibiting the ptosis induced in the mouse by tetrabenzain; the ED_{50} is of the order of 1 mg/kg. This test is used as a standard for antidepressant properties (International Journal of Neuropharmacology 8, 73 (1968)). 55

The compounds according to the invention have also shown an exceptionally favourable activity in the test for reserpine antagonism, namely the reversal of the hypothermic effect caused by reserpine by means of a substance with an antidepressant activity. It has also been found that they inhibit the re-absorption of serotonin and adrenalin into the neurones. 60

The new compounds are particularly useful in that they have a different structure from the antidepressants known up to now; they are equivalent or superior to known commercially available products in their activity but have a lower toxicity. 65

Particular mention should be made of compounds of general formula I wherein R₁ represents a bromine atom in the p-position, X represents an oxygen atom or the amino group, R₂ represents hydrogen and R₃ and R₄ represent methyl groups. Particular mention should be made of the compounds:

- 5 2-(p-bromophenyl)-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine, 5
2-(p-bromophenyl)-2-(pyridyl-(2)-amino)-N,N-dimethyl-ethylamine,
2-(p-bromophenyl)-2-pyridyl-(3)-amino)-N,N-dimethyl-ethylamine.

- According to a yet further feature of the present invention there are provided pharmaceutical compositions comprising as active ingredient, at least one compound of formula I as hereinbefore defined or a physiologically compatible acid addition salt thereof in association with a pharmaceutical carrier or excipient. 10

- For pharmaceutical administration the compounds of general formula I and their physiologically compatible acid addition salts may be incorporated into the conventional preparations in either solid or liquid form, optionally in combination with other active ingredients. The compositions may, for example, be presented in a form suitable for oral administration. 15
Preferred forms include, for example tablets and coated tablets

- The active ingredient may be incorporated in excipients customarily employed in pharmaceutical compositions such as, for example, talc, gum arabic, lactose, starch, magnesium stearate, cocoa butter, aqueous or non-aqueous vehicles, fatty substances of animal or vegetable origin, 20 paraffin derivatives, glycols, various wetting, dispersing or emulsifying agents and/or preservatives,

Advantageously the compositions may be formulated as dosage units, each unit being adapted to supply a fixed dose of active ingredient.

- The following non-limiting examples serve to illustrate the present invention. 25

Example 1

2-p-Bromophenyl-2-(pyridyl-(3)-oxy)-N,N-dimethyl-ethylamine dihydrochloride

- 29.9 g of 2-chloro-2-p-bromophenyl-N,N-dimethyl-ethylamine hydrochloride (0.1 mol) and 14.3 g of 3-hydroxypyridine (0.15 mol) are refluxed for 16 hours in 150 ml of 25% sodium hydroxide solution and 150 ml of toluene with 0.5g of benzyltriphenyl phosphonium chloride. 30
After the solvent has been distilled off from the organic phase a residue is left which is dissolved in alcohol. The dihydrochloride of the title compound is obtained with hydrochloric acid. For purification, the title compound is treated in cyclohexane with active charcoal and kieselguhr. The yield of dihydrochloride is 20 g (44% of theory).
35 Melting point 128–129°C (ethanol). 35

Example 2

2-p-Bromophenyl-2-(pyridyl-(2)-amino)-N,N-dimethyl-ethylamine-dioxalate

- 24.5 g of 2-p-bromophenyl-N,N-dimethylethanolamine (0.1 mol) and 11 g of 2-aminopyridine 40 (0.11 mol) are heated to 60°C for 30 minutes in 50 ml of methanesulphonic acid. Whilst still warm, the reaction mixture is poured on to ice, made alkaline with ammonia and extracted with ethyl acetate. The product is purified by column chromatography over silica gel/methylene chloride-ethyl acetate-methanol. The crystalline dioxalate of the title compound is obtained by the addition of alcoholic oxalic acid.
45 The yield is 21 g (42% of theory). Melting point 170–171°C. 45

The following compounds of general formula I were obtained using the process described hereinbefore:

5	Ex- ample No.	R ₁	R ₂	R ₃	R ₄	R ₅	X	Pyr- idyl bond	Mp°C	5
	3	4-Br	H	H	H	CH ₃	NH	2	181-182	
	4	H	H	H	CH ₃	CH ₃	O	2	161-162	
	5	H	H	H	CH ₃	CH ₃	O	3	151-152	
10	6	H	H	H	CH ₃	CH ₃	NH	2	129-130	10
	7	H	H	H	CH ₃	CH ₃	NH	3	123-125	
	8	4-Br	H	2-Cl	CH ₃	CH ₃	O	3	135-136	
	9	4-Br	H	5-Cl	CH ₃	CH ₃	O	2	234-235	
	10	4-Br	H	6-Cl	CH ₃	CH ₃	O	2	193-194	
15	11	H	H	4-CH ₃	CH ₃	CH ₃	NH	2	139-140	15
	12	4-Br	H	5-Cl	CH ₃	CH ₃	NH	2	134-136	
	13	4-Br	H	5-CH ₃	CH ₃	CH ₃	NH	2	103-104	
	14	4-Br	H	6-CH ₃	CH ₃	CH ₃	NH	2	135-136	
	15	4-Br	H	H	CH ₃	CH ₃	NH	3	144-145	
20	16	4-Br	H	H	CH ₃	CH ₃	O	2	162-163	20
	17	4-Br	H	H	CH ₃	CH ₃	S	2	190-191	
	18	4-Br	H	H	Morpholine		O	3	95-97	
	19	4-Br	H	H	Morpholine		NH	2	194-195	
	20	4-Br	H	H	Pyrrolidine		O	3	152-153	
25	21	4-Br	H	H	Pyrrolidine		NH	2	117-119	25
	22	4-Br	H	H	CH ₃	CH ₃	O	4	112 (decomp)	
	23	4-Cl	H	H	CH ₃	CH ₃	O	3	126-129 (decomp)	
	24	3-Cl	4-Cl	H	CH ₃	CH ₃	O	3	175-176	
30	25	4-OCH ₃	H	H	CH ₃	CH ₃	O	3	103-105	30
	26	3-Cl	4-Cl	H	CH ₃	CH ₃	NH	3	152-153	
	27	3-OCH ₃	H	H	H	H	O	3	222-223	
	28	4-CH ₃	H	H	CH ₃	CH ₃	O	3	180-181	
	29	4-Br	H	4-CH ₃	CH ₃	CH ₃	NH	2	219-220	
35										35

Examples of Formulation

a) Coated tablets

1 tablet core contains:

40	Active substance according to the invention	25.0 mg	40
	Lactose	50.0 mg	
	Corn starch	22.0 mg	
	Gelatine	2.0 mg	
45	Magnesium stearate	1.0 mg	45
		<u>100.0 mg</u>	

Preparation:

50	The mixture of active substance, lactose and corn starch is granulated with a 10% aqueous gelatine solution through a screen with a 1 mm mesh, then dried at 40°C and again passed through a screen. The granulate thus obtained is mixed with a magnesium stearate and compressed. The cores thus obtained are coated in the usual way with a coating which is applied using an aqueous suspension of sugar, titanium dioxide, talc and gum arabic. The	50
55	finished coated tablets are polished with beeswax. Final weight of coated tablet: 200 mg	55

b) *Tablets*

Active substance according to the invention

10.0 mg

5 Lactose

40.0 mg

5

Corn starch

44.0 mg

Soluble starch

5.0 mg

Magnesium stearate

1.0 mg

10

100.0 mg

10

Method:

The active substance and magnesium stearate are granulated with an aqueous solution of the soluble starch, the granulate is dried and intimately mixed with lactose and corn starch. The mixture is then compressed to form tablets weighing 100 mg and each containing 10 mg of active substance.

15

15

c) *Suppositories*

1 Suppository contains:

20

Active substance according to the invention

10.0 mg

20

Suppository mass

1,690.0 mg

Method:

25

Using an immersion homogeniser, the finely powdered substance is stirred into the molten suppository mass which has been cooled to 40°C. At 35°C the mass is poured into slightly pre-cooled moulds.

25

d) *Ampoules (Injection solutions)*

30

Composition:

Active substance according to the invention

5.0 parts by weight

Sodium pyrosulphite

1.0 parts by weight

Disodium salt of ethylene-diamine tetraacetec acid

0.5 parts by weight

35

35

Sodium chloride

8.5 parts by weight

Doubly distilled water *ad*

1,000 parts by weight

Method

40

The active substance and excipients are dissolved in sufficient water and adjusted to the desired concentration with the required amount of water. The solution is filtered and transferred into 1 ml ampoules under aseptic conditions. Finally the ampoules are sterilised and sealed. Each ampoule contains 5.0 mg of active substance.

40

45 CLAIMS

45

1. Substituted 2-phenyl-2-(pyridyloxy)-ethylamines and isosteric compounds of general formula

50


50

55

55

60

60



wherein

R_1 and R_2 , which may be the same or different, each represents a hydrogen or halogen atom or a methyl, methoxy, amino or nitro group;

65

R_3 represents a hydrogen or halogen atom or a methyl group;

65

R_4 and R_5 , which may be the same or different, either each represents a hydrogen atom or an alkyl group with 1 or 2 carbon atoms or, together with the nitrogen atom to which they are attached they may form a pyrrolidino or morpholino ring, and
X represents O, NH or S;

5 and acid addition salts thereof. 5


2. 2-(p-Bromophenyl)-2-(pyridyl-(3-oxy)-N,N-dimethyl-ethylamine and acid addition salts thereof.

3. 2-(p-Bromophenyl)-2-(pyridyl-(2-amino)-N,N-dimethyl-ethylamine and acid addition salts thereof.


10 4. 2-(p-Bromophenyl)-2-(pyridyl-(3-amino)-N,N-dimethyl-ethylamine and acid addition salts thereof. 10

5. Physiologically compatible acid addition salts of compounds of general formula I as defined in claim 1.

15 6. Process for the preparation of compounds as claimed in claim 1 which comprises reacting a phenylethylamine of formula 15



(wherein R_1 , R_2 , R_4 and R_5 are as defined in claim 1 and Y represents a halogen atom or a hydroxy group) or an acid addition salt of this compound, with a pyridine derivative of formula



(wherein R_3 is as defined in claim 1 and Z represents a hydroxy, amino or mercapto group) followed, if desired, by conversion of the product thus obtained into an acid addition salt thereof.

35 7. Process as claimed in claim 6 wherein, in the compound for formula II, Y represents a hydroxy group and the reaction is effected in an acidic medium at elevated temperatures. 35

8. Process as claimed in claim 6 wherein, in the compound of formula II, Y represents a halogen atom and the reaction is effected in the presence of a base.

9. Process for the preparation of compounds as claimed in claim 1 wherein R_4 and/or R_5 represents an alkyl group with 1 or 2 carbon atoms which comprises alkylating a compound as claimed in claim 1 wherein R_4 and/or R_5 represents a hydrogen atom. 40

10. Process as claimed in claim 9 wherein alkylation is effected by means of a dialkyl sulphate or an alkyl halide.

11. Process for the preparation of compounds as claimed in claim 1 substantially as herein 45 described. 45

12. Process for the preparation of compounds as claimed in claim 1 substantially as herein described in any one of Examples 1-29.

13. Process for the preparation of compounds as claimed in claim 1 substantially as herein described in any one of Examples 1-26.

50 14. Compounds as claimed in claim 1 whenever prepared by a process as claimed in any one of claims 6 to 12. 50

15. Compounds of general formula I as claimed in claim 1 and physiologically compatible acid addition salts thereof for use as antidepressants.

16. Pharmaceutical compositions comprising, as active ingredient, at least one compound of formula I as defined in claim 1 or a physiologically compatible acid addition salt thereof in association with a pharmaceutical carrier or excipient. 55

17. Compositions as claimed in claim 16 in the form of dosage units.

18. Pharmaceutical compositions as claimed in claim 16 substantially as herein described.

19. Pharmaceutical compositions substantially as herein described in Formulation examples

60 (a) - (d). 60

20. A method of treating a patient suffering from or susceptible to depression which comprises administering to the said patient an effective amount of a compound of formula I as defined in claim 1 or a physiologically compatible acid addition salt thereof.

21. Each and every novel method, process, compound and composition herein disclosed.

Printed for Her Majesty's Stationery Office by Burgess & Son (Abingdon) Ltd.—1983.
Published at The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.